

Formation of ω Centauri by Tidal Stripping of a Dwarf Galaxy

Makoto Ideta

*Astronomical Data Analysis Center, National Astronomical Observatory of Japan
2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*

and

Junichiro Makino

*Department of Astronomy, University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

ABSTRACT

We have investigated whether or not a tidal stripping scenario can reproduce the observed surface-brightness profile of ω Centauri using N -body simulations. Assuming that the progenitor of ω Centauri is a dwarf elliptical galaxy, we model it with a King model with a core radius being the same as that of ω Centauri. A dark matter halo of the dwarf is not taken into account. We consider two different models of the Milky Way potential: a singular isothermal sphere and a three-component model. The progenitor dwarf is expressed as an N -body system, which orbits in the fixed Galactic potential. The dwarf lost more than 90 per cent of its mass during the first few pericenter passages. Thereafter, the mass remains practically constant. The final surface-density profile is in good agreement with the observational data on ω Centauri, if the pericenter distance of the orbit of the progenitor dwarf is around 500 pc. This value is within the error bar of the current proper motion data on ω Centauri and Galactic parameters. Although our simulation is limited to a King-like progenitor dwarf without a dark matter halo, it strongly suggests that the current density profile of ω Centauri is nicely reproduced by a tidal stripping scenario, in other words, that ω Centauri can plausibly be identified with a stripped dwarf elliptical.

Subject headings: Galaxy: formation — globular clusters: individual (NGC 5139)
— methods: N -body simulations

1. INTRODUCTION

ω Centauri (NGC 5139) is, to the best of our knowledge, the most massive globular cluster in the Milky Way, with a mass around $5 \times 10^6 M_\odot$ (Meylan et al. 1995). Furthermore, it shows peculiar chemical and dynamical features, such as the wide spread in metallicity distribution of member stars (Freeman & Norris 1981) and the difference in kinematics of metal-rich and metal-poor populations (Norris et al. 1997). Some member stars show an enhancement of s-process elements produced by AGB stars (e.g., Smith et al. 2000), which means that the potential well must have been deep enough to trap the gas ejected from AGB stars (Gnedin et al. 2002). A merger model (Icke & Alcaïno 1988) can explain the kinematics and metallicity spread, but has difficulty in accounting for the presence of s-process elements.

A tidal stripping scenario (Zinnecker et al. 1988; Freeman 1993) has been proposed to explain all observed features simultaneously. In this scenario, the cluster was born as a dwarf elliptical galaxy, which was much more massive than the current ω Centauri. As it sinks to the Galactic center through dynamical friction, it loses most of its mass through the tidal stripping by the gravitational potential of the Milky Way. However, if the central density is sufficiently high, the core of the dwarf can still survive as a bound system and become a relatively large globular cluster, something like ω Centauri. In this scenario, a relatively long star formation history is naturally explained because the potential well of the progenitor dwarf elliptical was initially much deeper than that of the present-day ω Centauri. In addition, since the progenitor dwarf could be formed through hierarchical merging events, signatures of past merging events can also be explained. Furthermore, recently, Martini & Ho (2004) reported that some of massive globular clusters in NGC 5128 may also be such tidally stripped dwarf ellipticals.

From the viewpoint of dynamics, whether or not the stripping scenario works depends on the following two questions. The first one is whether the dynamical friction can bring the progenitor dwarf to the current orbit of ω Centauri. The second one is whether the tidal stripping can actually produce the observed surface brightness of ω Centauri. The first question has been addressed by recent work (Zhao 2002; Tsuchiya, Dinescu, & Korchagin 2003; Bekki & Freeman 2003). However, the second question has not been studied so far, despite its significance to the validity of the stripping scenario.

In this paper, we have investigated whether a dwarf galaxy with the orbit similar to that of the present-day ω Centauri will evolve to have the spatial structure which agrees with that of ω Centauri using N -body simulations. In § 2, numerical models and method are described. Results are shown in § 3. A discussion is given in § 4.

2. MODELS AND METHOD

We consider the dynamical evolution of a dwarf elliptical in a fixed potential. In this paper, we neglect the effects of dynamical friction. This neglect should not affect results as far as a progenitor dwarf along the present-day orbit of ω Centauri is concerned, since our main goal is to see whether or not the tidal field of the Milky Way in the present-day orbit of ω Centauri can account for its density structure. In reality, dynamical friction must have played a certain role in bringing the progenitor dwarf to its current orbit, but that part of evolution is not in the scope of the present paper.

We use the data compiled by Meylan (1987) for the surface-brightness profile of ω Centauri. This profile is well-fitted (except in the outermost region) by a King model with a non-dimensional central potential $W_0 = 5.5$ and a core radius $R_c = 4.6$ pc. We choose the total mass of this King model to be identical to the mass suggested from observations, $5.0 \times 10^6 M_\odot$. Then, we set up the model of the progenitor dwarf to have the same core radius and central velocity dispersion, but with a much more extended halo. We choose a King model with $W_c = 12$ as a model for the dwarf. Since some dwarf elliptical galaxies can also be fitted by a King model, we believe this choice is justified. Tsuchiya, Korchagin, & Dinescu (2004) have found that a King-like dwarf elliptical cannot evolve to ω Centauri on the basis of numerical simulations of King- and Hernquist-like dwarf galaxies embedded in a live Milky Way potential. This point will be discussed in § 4.1.

The dwarf elliptical is represented by $2^{20} \simeq 1$ million particles of equal mass. The initial total mass is $1.3 \times 10^8 M_\odot$. We do not include a dark matter halo of the dwarf, which would be more extended than the distribution of visible stars. Since the extended halo would be stripped out anyway, we believe the presence of a dark matter halo does not change results significantly. In most of our simulations, the final cluster still had more than 50,000 particles, with the result that the relaxation time is much longer than the duration of the simulation.

We use two different models for the gravitational potential of the Milky Way, a singular isothermal sphere (run std) and a bulge-disk-halo model (run mwd). The circular velocity of the std potential is 220 km sec^{-1} . The mwd potential is almost the same as the model used in Johnston, Spergel, & Hernquist (1995). This model consists of three components: a Miyamoto-Nagai (1975) disk, a Hernquist (1990) bulge, and a logarithmic halo potential. The only difference between the model of Johnston et al. (1995) and ours is the core radius of the logarithmic potential. We set the core radius to be 14.0 kpc, while Johnston et al. (1995) set it to be 12.0 kpc, so that the circular velocity at the solar circle radius, i.e., 8.0 kpc, is 220 km sec^{-1} . The potentials and parameters used in simulations are listed in Table 1. The circular speed curves of these models are plotted in Figure 1.

We use the following kinematical data on ω Centauri: the distance from the Sun is 4.9 kpc, the proper motions are $(\mu_\alpha \cos \delta, \mu_\delta) = (-5.08 \pm 0.35, -3.57 \pm 0.34)$ mas yr $^{-1}$, and the radial velocity is 232.2 ± 0.7 km sec $^{-1}$ (Dinescu et al. 1999). For the most probable values of the proper motions and the radial velocity in the isothermal model (run std), the pericenter and apocenter distances are 1.0 kpc and 6.4 kpc, respectively. We made two other runs, in which we expand the initial dwarf by a factor of 1.5 (run r15) and 2 (run r20), to see the effect of the change in the tidal force at the pericenter. For the singular isothermal model, the only characteristic scale in the experiment is the core radius of the cluster. A linear expansion of the cluster by a factor of 1.5 or 2 can therefore equally well correspond to a reduction in the pericenter distance of the unmodified cluster by a factor of $1.5^{3/2} \simeq 2$ (run r15) or $2^{3/2} \simeq 3$ (run r20), which corresponds to ~ 600 pc and ~ 400 pc, respectively.

We use a hierarchical tree algorithm (Barnes & Hut 1986) on the GRAPE-6 hardware (Makino et al. 2003) with an opening angle being 0.5. We employ a Plummer softening of 0.5 pc, which is roughly one-tenth of the observed core radius of ω Centauri, 4.6 pc. The equations of motion are integrated using a leap-frog method with a constant time-step of $\Delta t = 1.7 \times 10^4$ yr. This time-step is comparable to the time required for a particle with the maximum circular velocity around ω Centauri to cover the softening length. The total energy was conserved to better than 0.005 per cent in all simulations.

3. RESULTS

Figure 2 shows snapshot images from run std, along with the orbit of the cluster center for the periods of 0.1 Gyr before and after the time for the snapshot (whenever the orbit is available). The cluster center is defined as the position of the particle with the minimum potential energy for the N -body particles. We include all particles to calculate the potential. Removing unbound particles did not affect the result, even after more than 90 per cent of the particles were stripped. We can see that the stripped stars remain close to the cluster orbit, and form numerous ripple structures.

Figure 3 shows the time evolution of the cluster mass. Here, we define the cluster mass simply as a mass within $2r_t$, where r_t is the tidal radius at the apocentric position expressed as

$$r_t = \left(\frac{m_c}{2M_G} \right)^{1/3} R_G. \quad (1)$$

Here R_G is the distance from the Galactic center to the cluster, M_G is the Galaxy mass within R_G , and m_c is the cluster mass. We assume that the stars are stripped from the cluster if their distances from the cluster center are larger than two apocentric tidal radii (e.g., Baumgardt

& Makino 2003). For the three-component model, we have determined the cluster mass in the same way. The cluster mass exhibits periodic sudden drops, which correspond to the pericentric passage. Thus, almost all the mass loss occurred at the pericentric passage. Since the cluster is expanded in runs r15 and r20, the mass loss is larger for these runs than that for run std. There is very little difference between the result of run std and that of mwd, suggesting that the disk does not contribute significantly to the tidal force. This result is of course not surprising, because the disk gravity is small at a pericenter distance of 1 kpc.

Figure 4 shows the final surface-density profiles for all runs. We assume a constant mass-to-light ratio, since the estimated relaxation time of ω Centauri is too long for the mass segregation to be visible (e.g., Giersz & Heggie 2003). Clearly, the result of run std does not agree with observations. Replacing the Galactic potential with a more realistic one (run mwd) does not improve the situation. On the other hand, changing the pericenter distance has a drastic effect, and the agreement of run r20 with observations is very good. The result of r15 falls between std and r20.

We conclude that a tidal stripping scenario involving a King-like progenitor without a dark matter halo can reproduce the observed luminosity profile quite well, provided that the pericentric distance of ω Centauri is around 400 pc.

4. DISCUSSION

4.1. Comparison with Previous Work

Our conclusion that a King-like dwarf can evolve to ω Centauri apparently contradicts with the result by Tsuchiya et al. (2004) that final clusters started from King-like dwarf ellipticals are still more massive than $10^8 M_{\odot}$. This difference is mainly due to the high velocity dispersion of the initial dwarf models in their simulations. The central velocity dispersion in their model K5 reaches $\sim 55 \text{ km sec}^{-1}$, while we set it to be 22 km sec^{-1} . This means that the cluster mass inside the tidal radius in their model K5 is $2.5^3 \simeq 16$ times larger than the mass in our model, even if the pericenter distance is assumed to be equal. In terms of the fraction of total mass lost, their results are actually quite consistent with ours.

4.2. Pericenter Distance

We have shown that the observed surface-brightness profile of ω Centauri is nicely reproduced by the tidal stripping scenario if the pericenter distance of the cluster is roughly 400 pc.

Here, we discuss whether such a small pericenter distance is compatible with observations.

We have adopted the following error estimates for observational data. For the errors in the proper motion and the radial velocity, we have used the data by Dinescu et al. (1999): $(U, V, W) = (-64 \pm 11, -254 \pm 9, 4 \pm 10)$ km sec⁻¹. Here, (U, V, W) is the relative velocity of ω Centauri to the local standard of rest. The positive directions of the U , V , and W components are, respectively, outward from the Galactic center, toward the Galactic rotation, and toward the Galactic north pole. For the Galactic parameters, Reid et al. (1999) estimated the ratio of the circular speed to the solar circle radius, Θ_0/R_0 , on the basis of the proper motion study of Sagittarius A* by using VLBA, and concluded that the circular speed is 219 ± 20 km sec⁻¹, assuming that the solar circle radius is $R_0 = 8.0$ kpc. For the error in the circular speed at the solar circle, we have adopted the value of 220 ± 20 km sec⁻¹. Since the solar circle radius would have an error comparable to that in the circular speed, we have used 8.0 ± 1.0 kpc for the solar circle radius.

We calculated the variations of the pericentric distance due to errors in the orbital velocity of ω Centauri, the solar circle radius, and the circular speed at the solar circle. The results are summarized in Table 2. The error in the circular speed has a strong effect on the pericenter distance, since the orbit of ω Centauri is highly eccentric. A 10 per cent variation in the circular speed can lead to 50 per cent change in the pericenter distance. The error in the V component also has such large magnification. Thus, our result that the pericenter distance must be around 400 pc is compatible with observations.

We set the total mass of ω Centauri to $5.0 \times 10^6 M_\odot$, following the estimate by Meylan et al. (1995). However, since this mass estimate is based on a multi-mass King model, the total mass is probably overestimated (see Baumgardt et al. 2003). Indeed, in our best-fit King model, the central velocity dispersion reaches 22 km sec⁻¹, while the observed value is around 17 km sec⁻¹ (Merritt, Meylan, & Mayor 1997). For the observed value of the central velocity dispersion, the total mass is around $3 \times 10^6 M_\odot$. This change of the cluster mass reduces the tidal radius by about 15 per cent ($r_t \propto m_c^{1/3}$). In other words, to yield the observed tidal radius with a reduced cluster mass, the periastron distance in the model would have to be increased by 30 per cent [from equation (1), $R_G \propto m_c^{-1/2}$]. Therefore, if we use this revised estimate of the mass of ω Centauri, the most likely value of the pericenter distance is around 500 pc.

4.3. Long-Term Orbit Evolution in the Three-Component Model

In section 3, we have found that the quantitative result remains unchanged when we change the Galactic potential model from the singular isothermal sphere to the more realistic three-component model. However, since we integrated our model only for 0.88 Gyr, we could have missed some important contributions of the non-sphericity of the potential.

We integrated the orbit of a point-mass cluster in the three-component potential for $\sim 10^{13.5}$ yr. Here, such a long integration time is required since chaotic diffusion time could be much longer than a Hubble time. We have found that the minimum pericentric radius is $\simeq 0.9$ kpc, although the orbit is chaotic. Consequently, the effect of chaotic diffusion is negligible.

4.4. Globular Clusters as Probes of Galactic Structure

We have found that the structure of ω Centauri is nicely explained by the tidal stripping scenario, for the orbital parameters within the observational errors. The pericenter distance is the most important parameter that determines the structure of the simulated ω Centauri. In other words, we can put a fairly tight constraint on the pericenter distance of ω Centauri from its internal structure. This constraint can be converted to constraints on the Galactic potential, such as the circular speed, if we have high-accuracy data for the proper and radial motions of ω Centauri.

We can probably apply a similar technique to other tidally-limited globular clusters, though in this case the modeling is somewhat more complex since we cannot ignore thermal relaxation. Clusters in highly eccentric orbits are most useful, since a small change in the Galactic parameters results in a large change in the pericenter distance. Thus, the dynamical simulation of the evolution of globular clusters in a Galactic tidal field, combined with high-accuracy proper motion data which will be available via next-generation astrometry projects, will provide us with a unique tool to probe the structure of the Milky Way.

We thank the anonymous referee for useful comments that helped us to improve the clarity of the paper. We are grateful to Prof. S. Hozumi for his critical reading of the manuscript. All simulations were run at the University of Tokyo. Some part of data analysis was made on workstations at Astronomical Data Analysis Center, National Astronomical Observatory of Japan (ADAC/NAOJ).

REFERENCES

- Barnes, J., & Hut, P. 1986, *Nature*, 324, 446
- Baumgardt, H., & Makino, J. 2003, *MNRAS*, 340, 227
- Baumgardt, H., Makino, J., Hut, P., McMillan, S., & Portegies Zwart, S. 2003, *ApJ*, 589, L25
- Bekki, K., & Freeman, K. C. 2003, *MNRAS*, 346, L11
- Dinescu, D. I., van Altena, W. F., Girard, T. M., & López, C. E. 1999, *AJ*, 117, 277
- Freeman, K. C. 1993, in *ASP Conf. Ser. 48, The Globular Cluster-Galaxy Connection*, ed. G. H. Smith, & J. P. Brodie (San Francisco: ASP), 608
- Freeman, K. C., & Norris, J. 1981, *ARA&A*, 19, 319
- Giersz, M., & Heggie, D. C. 2003, *MNRAS*, 339, 486
- Gnedin, O. Y., Zhao, H., Pringle, J. E., Fall, S. M., Livio, M., & Meylan, G. 2002, *ApJ*, 568, L23
- Hernquist, L. 1990, *ApJ*, 356, 359
- Icke, V., & Alcaïno, G. 1988, *A&A*, 204, 115
- Johnston, K. V., Spergel, D. N., & Hernquist, L. 1995, *ApJ*, 451, 598
- Makino, J., Fukushige, T., Koga, M., & Namura, K. 2003, *PASJ*, 55, 1163
- Martini, P., & Ho, L. C. 2004, *ApJ*, 610, 233
- Merritt, D., Meylan, G., & Mayor, M. 1997, *AJ*, 114, 1074
- Meylan, G. 1987, *A&A*, 184, 144
- Meylan, G., Mayor, M., Duquennoy, A., & Dubath, P. 1995, *A&A*, 303, 761
- Miyamoto, M., & Nagai, R. 1975, *PASJ*, 27, 533
- Norris, J. E., Freeman, K. C., Mayor, M., & Seitzer, P. 1997, *ApJ*, 487, L187
- Reid, M. J., Readhead, A. C. S., Vermeulen, R. C., & Treuhaft, R. N. 1999, *ApJ*, 524, 816

- Smith, V. V., Suntzeff, N.B., Cunha, K., Gallino, R., Busso, M., Lambert, D. L., & Straniero, O. 2000, *AJ*, 119, 1239
- Tsuchiya, T., Dinescu, D. I., & Korchagin, V. I. 2003, *ApJ*, 589, L29
- Tsuchiya, T., Korchagin, V. I., & Dinescu, D. I. 2004, *MNRAS*, 350, 1141
- Zhao, H. S. 2002, in *ASP Conf. Ser. 265, ω Centauri: A Unique Window into Astrophysics*, ed. F. van Leeuwen, J. D. Hughes, & G. Piotto (San Francisco: ASP), 391
- Zinnecker, H., Keable, C. J., Dunlop, J. S., Cannon, R. D., & Griffiths, W. K. 1988, in *IAU Symp. 126, Globular Cluster Systems in Galaxies*, ed. J. E. Grindlay & A. G. D. Philip (Dordrecht: Kluwer), 603

Table 1. The Milky Way models.

isothermal sphere	three component
$\Phi = v_c^2 \ln r$	$\Phi_d = -\frac{GM_d}{\sqrt{R^2 + (a + \sqrt{z^2 + b^2})^2}}$
$v_c = 220 \text{ km sec}^{-1}$	$M_d = 1.0 \times 10^{11} M_\odot, \quad a = 6.5 \text{ kpc}, \quad b = 0.26 \text{ kpc}$
	$\Phi_b = -\frac{GM_b}{r+c}$
	$M_b = 3.4 \times 10^{10} M_\odot, \quad c = 0.7 \text{ kpc}$
	$\Phi_h = v_h^2 \ln (r^2 + d^2)$
	$v_h = 128 \text{ km sec}^{-1}, \quad d = 14 \text{ kpc}$

Table 2. The pericenter distance of ω Centauri in the isothermal sphere model.

observable	value	pericenter
velocity (U, V, W)	$(-64 \pm 11, -254 \pm 9, 4 \pm 10) \text{ km sec}^{-1}$	$1.0 \pm 0.3 \text{ kpc}$
solar circle radius	7.0 kpc	0.9 kpc
	9.0 kpc	1.0 kpc
circular speed	240 km sec^{-1}	0.6 kpc
	200 km sec^{-1}	1.5 kpc




Fig. 1.— Circular speed curves of our Galactic models. The dashed, dash-dotted, and dotted curves give the contributions of the bulge, disk, and halo, respectively, while the solid curve gives the total circular speed in the three-component model. The horizontal dotted line at 220 km sec^{-1} is the rotation curve of the isothermal model.




Fig. 2.— Snapshot images from run std. Bound particles are drawn in black, while unbound ones are in gray. Simulation time is shown at the top-right corner of each box. The orbit of the cluster center for periods of 0.1 Gyr before and after each simulation time is also drawn in the solid line.




Fig. 3.— Time evolution of the cluster mass for all runs. The solid, dotted, dash-dotted and dashed curves represent the results of runs std, mwd, r15, and r20, respectively.




Fig. 4.— Final surface-density (brightness) profiles of the clusters for all runs. The curves have the same meanings as in figure 3. The open circles with error bars show the surface-density profile calculated from the observed surface brightness on the assumption of a constant mass-to-light ratio. The thin solid line corresponds to the initial model.